

# Coral communities in extreme environmental conditions in the Northern Territory, Australia

Lawrance W. Ferns

Biodiversity Division, Department of Environment, Land, Water and Planning,  
8 Nicholson Street, East Melbourne, Victoria 3002, Australia  
Email: [lawrance.fern@delwp.vic.gov.au](mailto:lawrance.fern@delwp.vic.gov.au)

## Abstract

An extensive intertidal reef flat in the macro-tidal marine waters of the Northern Territory was chosen to investigate species composition and zonation persisting under extreme environmental conditions. Thirty-six visual belt transects were used to quantify scleractinian corals, benthic algae and other sessile invertebrates which varied in vertical and horizontal space. Thirty-four coral species were identified. Most species were represented by the family Merulinidae, with lifeform characteristics typical of species specialised in environmental tolerance to high sedimentation, turbidity and temperature (i.e. massive, sub-massive and encrusting growth forms with convex and steep sided morphologies, thick skeletal tissue and large polyps). Whilst the combination of environmental and ecological characteristics of this reef flat community can be viewed as distinctive to the Darwin region, a number of similarities can be compared to reef communities reported in extreme environments of the Arabian Gulf, Red Sea and other regions of tropical northern Australia.

---

## Introduction

Coral communities which presently persist in extreme environmental conditions are of contemporary interest towards understanding species resilience and potential for adaptation to climate change (Hughes *et al.* 2003; Bauman *et al.* 2011, 2013a, 2013b; Dandan *et al.* 2015). Predictions for climate change stressors for coral communities include warmer sea temperature and changes in extreme episodic events such as heavy rainfall, storms and possible sediment and nutrient debouching from rivers and run-off (Gilmour *et al.* 2006). Such environmental perturbations are known to be associated with mass coral bleaching and sudden die off (Glynn 1993; Depczynski *et al.* 2013). Coral species of the coastal waters in the vicinity of Darwin Harbour, Northern Territory (Wolstenholme *et al.* 1997), the Kimberley coast, northwestern Australia (Dandan *et al.* 2015; Schoepf *et al.* 2015) and the Arabian Gulf (Sheppard & Sheppard 1991; Coles 1997; Coles 2003; Riegl 1999; Bauman *et al.* 2011; Riegl & Purkis 2012) survive in harsh conditions and offer insight to species with high climatic tolerances and adaptation.

Climatic trends in this region of Australia are changing. Since 1950, the Northern Territory average rainfall has risen 35.7 mm per decade during November–April and

---

fallen 0.4 mm per decade during May–October. From 1910 to 2003, the intensity of heavy daily rainfall gradually rose by 10% (Hennessy *et al.* 2004). Water temperatures at the shoreline may reach over 36°C at high tide, with tide pools and standing water bodies reaching over 43°C at low tide which is similar to the high temperature fluctuations in the Arabian Gulf, and well above temperature ranges corals traditionally considered limiting to coral survival (Coles 1997). Sea surface temperatures north of Australia have been at record-breaking highs. In 2010, temperatures north of Australia broke previous records by large margins and were also above average during the 2011–2012 La Niña event (Australian Bureau of Meteorology 2013).

An extensive intertidal reef flat community off the north-eastern shoreline of Darwin Harbour was chosen to investigate coral zonation on a macro-tidal shoreline. The marine waters of Darwin Harbour are subjected to daily and seasonal fluctuations in sea surface temperature, light availability and extreme levels of sedimentation and turbidity which provides insight to northern Australian coral species that have already adapted to extreme climatic conditions.

## Methods


### Study site

Nightcliff Reef (Fig. 1) was surveyed at low spring tide between September and October 1994. Darwin Harbour is a ria coast formed by the post-glacial marine flooding of a dissected plateau. Subsequent sedimentary infill has resulted in the formation of numerous embayments, islands and extensive mangrove-vegetated tidal flats (Semeniuk 1985). The dominant lithological type is ferricrete laterite with phyllite/siltstone and this is reflected by the presence of conspicuous medium to coarse grained lateritic pebbles of the upper beach sediments. The coral community at Nightcliff exists as a veneer reef by colonising hard substrates of rock and consolidated materials without accreting substantial calcium carbonate substrata (Hooper 1987; Mitchie 1987). The community extends seaward to a distance of approximately 500 m from the shoreline at low spring tide.

Tides in Darwin Harbour exhibit semidiurnal inequality with a spring tidal range is in the vicinity of 0.1–7.8 m. Tidal currents are very strong ranging from 0.25–1.4 m/s and recording as high as 2 m/s (Semeniuk 1985). The concentration of rivers, streams and creeks with accompanying discharge, and the strong tidal movement of the system, account for excessive sediment mixing and high turbidity (Michie 1987). Nightcliff Reef is fully emersed for 2–3 hours during low spring tides that are less than 1 m. Low spring tides occur between 1100–1600 hrs in the period September to March (wet season) and between 2300–0400 hrs in the period April to August (dry season).

### Stratification

A topographic contour map of the reef was generated from random spot heights measured using a digital theodolite. All heights were referenced from a survey datum



**Fig. 1.** Location of Nightcliff Reef off the north-eastern foreshore of Darwin Harbour.

belt transects that were haphazardly placed in each of the upper, middle, and lower reef flat vertical zones. The sampling was repeated in three localities (north, central and south locations) to account for any variations in substrata micro-topography, sediment deposition and taxa composition that may occur horizontally across the reef relative to the shoreline. In total, 36 belt transects were surveyed. Contiguous belt transect was chosen as the sampling unit to improve the recording of small and less abundant species, and representation of microhabitats (e.g. large or small coral colonies, sand patches, tide pools) in any given zone and locality (Chiappone & Sullivan 1991; Sullivan & Chiappone 1992). Species abundances of all sessile taxa were recorded by visual estimates of percentage cover and numbers of individuals for each 1 m<sup>2</sup> quadrat. Lifeform attributes of coral colonies were recorded using categories of English *et al.* (1994). When a particular species was encountered that could not be identified in the field, a sample of the species was collected for identification in the laboratory and also compared to collection specimens held at the Northern Territory Museum. Original taxonomic identifications for scleractinian corals followed that of Veron and Pichon (1976, 1980, 1982); Veron *et al.* (1977); Veron and Wallace (1985) and Veron (1986). Algal identifications followed that of Jaasund (1976); Cribb (1983); Cribb and Cribb (1985); Lawson and John (1987) and Price and Scott (1992). Updates to recent taxonomic revisions follow Wynne (2011); Budd *et al.* (2012); Guiry and Guiry (2013); Huang *et al.* (2014) and WoRMS Editorial Board (2015). All data presented in the text and figures are the arithmetic mean.

## Results and Discussion

A total of 75 sessile species comprising scleractinian corals, algae, and sessile invertebrates were identified in the 36 transects sampled across the upper, middle and lower reef flat zones. 34 species of scleractinian corals from eight families were recorded (Table 1). The majority of species were members of the family Merulinidae, represented by 18 species (or 53% of all corals), followed by the Lobophyllidae and Poritidae, each represented by 4 species (23% of all corals). All remaining 5 families were represented by 1 or 2

point adjacent to Nightcliff Pier, and were converted to metres above the lowest astronomical tide (i.e. 0 m). The contour map was used to stratify the reef into three main vertical zones: (1) the upper reef flat which occupied a vertical height between 1.4–1.8 m; (2) the middle reef flat between 0.9–1.4 m; and (3) the lower reef flat between 0.3–0.9 m.

## Sampling

Sampling was conducted using four replicate 1 m x 20 m contiguous

species (24% of all corals). The more abundant corals, all with greater than 2% in mean percentage cover pooled for all 36 transects, were *Platygyra sinensis* (Fig. 2), *Porites* cf *nigrescens*, *Astrea curta*, *Coelastrea aspera* and *Goniastrea retiformis*. Secondary dominant corals (1–2%) were *Porites lutea*, *Lobophyllia hemprichii*, *Platygyra daedalea*, *Galaxea astreata*, *Favites*



**Fig. 2.** Massive colony of *Platygyra sinensis* at Nightcliff Reef, Darwin Harbour. This particular species and *Coelastrea aspera* have been sighted attaining colony sizes of up to 1.8 m in height. (Lawrance Ferns)

**Table 1.** Mean percentage cover of scleractinian corals, algae, other sessile invertebrates and physical substrates at Nightcliff Reef. Lifeform categories M = massive, S = submassive, E = encrusting, D/C = digitate/corymbose, R = ramose, T = turf, BF = bladed foliose, F = foliose, EC = erect coralline.

Family	Species	Lifeform	Upper Reef Flat (≥1.4–1.8 m)			Middle Reef Flat (≥0.9–1.4 m)			Lower Reef Flat (0.3–0.9 m)		
			north	central	south	north	central	south	north	central	south
CORALS											
Merulinidae	<i>Platygyra sinensis</i>	M		0.01		0.21		0.21	19.60	4.13	1.48
Poritidae	<i>Porites cf nigrescens</i>	S/R				8.10			4.34	5.76	3.75
Merulinidae	<i>Astrea curta</i>	M		0.06		2.85	0.15	1.66	3.93	3.25	2.35
Merulinidae	<i>Coelastrea aspera</i>	M							2.20	2.84	0.60
Merulinidae	<i>Goniastrea retiformis</i>	M	0.03	0.18	0.24	0.20	0.04	0.08	2.80	0.46	0.11
Poritidae	<i>Porites lutea</i>	M				0.50				1.36	
n/a	Dead standing coral	n/a				0.45			0.61	0.46	0.34
Lobophylliidae	<i>Lobophyllia hemprichii</i>	S							1.79		
Merulinidae	<i>Platygyra daedalea</i>	M				0.23			1.19	0.26	0.11
Euphylliidae	<i>Galaxea astrea</i>	E				0.13			1.24	0.25	
Merulinidae	<i>Favites abdita</i>	S				0.30	0.03		0.60	0.43	0.15
Merulinidae	<i>Dipastrea speciosa</i>	M	0.20	0.05		0.32		0.03	0.28	0.34	0.04
Merulinidae	<i>Dipastrea rotumana</i>	M				0.23			0.48	0.14	0.09
n/a	Juvenile corals	n/a	0.11	0.08		0.09		0.06	0.32	0.16	0.09
Merulinidae	<i>Dipastrea matthaii</i>	M				0.15			0.33	0.08	0.04
Merulinidae	<i>Dipastrea amicornum</i>	M	0.03			0.13			0.09	0.29	0.04
Merulinidae	<i>Cyphastrea serailia</i>	E	0.06	0.05		0.13	0.05		0.01	0.13	
Merulinidae	<i>Dipastrea favus</i>	M	0.12	0.04		0.17			0.03	0.05	
Acroporidae	<i>Acropora millepora</i>	D/C							0.34	0.05	
Merulinidae	<i>Dipastrea pallida</i>	M	0.01	0.10		0.16				0.04	
Poritidae	<i>Porites sp.l</i>	M				0.24		0.03			
Poritidae	<i>Goniopora cf. lobata</i>	S				0.04			0.14	0.09	
Lobophylliidae	<i>Symphyllia recta</i>	M				0.01			0.23		
Acroporidae	<i>Montipora encrusting</i>	E							0.13	0.06	
Dendrophylliidae	<i>Turbinaria mesenterina</i>	E				0.03			0.10		
Dendrophylliidae	<i>Turbinaria conspicua</i>	T							0.10		
Merulinidae	<i>Merulina ampliata</i>	E							0.10		
Scleractinia incertae sedis	<i>Leptastrea purpurea</i>	E								0.09	
Lobophylliidae	<i>Echinophyllia aspera</i>	E							0.09		
Scleractinia incertae sedis	<i>Leptastrea transversa</i>	E	0.01		0.01	0.02	0.01		0.00	0.03	
Merulinidae	<i>Cyphastrea microphthalma</i>	E							0.08		
Merulinidae	<i>Goniastrea pectinata</i>	E				0.03			0.04		

Continued on next page



Continued from previous page

Family	Species	Lifeform	Upper Reef Flat (>1.4–1.8 m)			Middle Reef Flat (>0.9–1.4 m)			Lower Reef Flat (0.3–0.9 m)		
			north	central	south	north	central	south	north	central	south
Merulinidae	<i>Cyphastrea chalcidicum</i>	E								0.05	
Psammocoridae	<i>Psammocora configua</i>	S				0.03					
Merulinidae	<i>Goniastrea favulus</i>	M							0.03		
Lobophylliidae	<i>Moseleya latistellata</i>	E							0.01		
Subtotal: Corals			0.57	0.57	0.25	14.75	0.28	2.07	41.23	20.80	9.19
ALGAE											
Sargassaceae	<i>Sargassaceae Holdfasts</i>	BF	4.01	0.43			1.81	0.08			
Dictyotaceae	<i>Padina australis</i>	F	12.55	20.39	6.84	0.03	0.63	0.15			0.03
Galaxauraceae	<i>Tricleocarpa fragilis</i>	T	0.16	0.61	0.69		0.23	0.01			
Dictyotaceae	<i>Dictyopteris sp.</i>	BF	0.09	1.16			0.01				
Corallinaceae	<i>Amphirna fragilissima</i>	T					2.72	0.23			
Sargassaceae	<i>Sargassopsis decurrens (Sighted only)</i>	BF									
Rhizophyllidaceae	<i>Portieria bornemannii</i>	T	1.21	0.24	0.05		0.30	0.20			
Dictyotaceae	<i>Spatoglossum asperum</i>	BF	0.32	1.24							
Halimeda	<i>Halimeda cf tuna</i>	EC	0.01		0.01		0.12				
Rhodomelaceae	<i>Palisada perforata</i>	T	0.11	0.21	1.21						
Rhodomelaceae	<i>Digenea simplex</i>	T									
Cystocloniaceae	<i>Hypnea spinella</i>	T	0.28		0.08						
Rhodomelaceae	<i>Acanthophora spicifera</i>	T	0.12	0.13	0.86		0.11				
Gracilariaceae	<i>Gracilaria salicornia</i>	T	0.28	0.33	0.61						
Halimeda	<i>Halimeda opuntia</i>	EC	0.13	0.21	0.01		0.27	0.11			
Anadyomenaceae	<i>Anadyomene plicata</i>	EC	0.22	0.21	0.07		0.26	0.33			
Cystocloniaceae	<i>Hypnea valentiae</i>	T	0.24	0.10	0.45						
Scytosiphonaceae	<i>Hydroclathrus clathratus</i>	T	0.08	0.51	0.13						
Caulerpaceae	<i>Caulerpa lentillifera</i>	F			0.01		0.08				
Rhodomelaceae	<i>Laurencia majuscula</i>	T		0.04	0.09						
Rhodomelaceae	<i>Laurencia intricata</i>	T		0.12	0.05		0.17				
Rhodomelaceae	<i>Tolytiocladia sp.</i>	T	0.01	0.01	0.22		0.09				
Callithamniaceae	<i>Crouania attenuata</i>	T	0.01	0.03	0.29						
Caulerpaceae	<i>Caulerpa racemosa</i>	F					0.03				
Galaxauraceae	<i>Dichotomaria obtusata</i>	T									
Caulerpaceae	<i>Caulerpa serrulata</i>	F	0.04								

Continued on next page

Continued from previous page

Family	Species	Lifeform	Upper Reef Flat (>1.4–1.8 m)			Middle Reef Flat (>0.9–1.4 m)			Lower Reef Flat (0.3–0.9 m)		
			north	central	south	north	central	south	north	central	south
Gelidiellaceae	<i>Gelidiella acerosa</i>	T	0.08	0.05	0.03						
Rhodomelaceae	<i>Laurencia obtusa</i>	T	0.05		0.04						
Cystocloniaceae	<i>Hypnea pannosa</i>	T				0.02			0.03		0.02
Cystocloniaceae	<i>Hypnea cornuta</i>	T			0.02						
Ulvaceae	<i>Ulva intestinalis</i>	F	0.03		0.02						
Champiaceae	<i>Champia parvula</i>	T					0.04				
Pithophoraceae	<i>Dictyosphaeria cavernosa</i>	F		0.04							
Cystocloniaceae	<i>Hypnea cf hamulosa</i>	T			0.03						
Solieriaceae	<i>Sarcinoma filiforme</i>	T	0.01		0.01						
Dichotomosiphonaceae	<i>Arrainvillea erecta</i>	F						0.01			
Dasycladaceae	<i>Neomeris annulata</i>	F	0.01								
Dictyotaceae	<i>Dictyota bartayresiana</i> (Sighted Only)	BF									
Sub-total: Algae			20.05	26.06	11.82	0.05	6.87	1.12	0.03	0.05	
SESSILE INVERTEBRATES											
Sponges			0.25	0.28	0.03	17.58	2.63	0.09	0.48	0.08	0.04
Other fauna (ascidians, soft corals)				0.03		0.04				0.01	
BARE SUBSTRATES											
Coarse sand			2.84	11.79		57.40			5.26	3.46	4.06
Coral rubble						6.90			53.01	75.67	86.84
Muddy Rock			72.88	55.93	74.51		82.09	78.12			
Muddy Sand			3.54	5.48	13.41	3.31	8.15	18.59			
Total (% Cover)			100	100	100	100	100	100	100	100	100

*abdita*, *Dipastrea speciosa*, and *Dipastrea rotumana*. Corals with ‘massive’ and ‘sub-massive’ lifeforms were the most common growth morphologies with 20 species (or 59% of all corals). The other main lifeform was ‘encrusting’ comprising 12 species (35% total of all corals). Only one species with digitate and tabulate lifeform categories were recorded (6% of all corals).

Coral abundance and species richness varied between vertical zones and horizontal localities across the reef. Physical factors such sediment type, micro-topography, standing water, with the relative influence of each factor shifting between and within vertical zones and localities, is the likely cause of the observed variation (Table 1). Species cover and richness increased markedly seaward from the upper reef flat zone (1.4–1.8 m) to the lower reef flat zone (0.3–0.9 m) which has the highest coral abundance and species richness. This is consistent with similar studies that have found intertidal corals to be more successful below 1 m tidal height elevations and reaffirms the period of emersion time is an important factor determining the their upper vertical limit (Morrissey 1980;

Bull 1982). Coral abundance and species richness was significantly higher on the lower reef flat at the northern extent of the reef (41% mean cover, 26 species), and decreased towards the southern extent (9% mean cover, 13 species). This is attributed to a notable decrease in fine sediments at the northern extent, combined with a higher proportion of coral rock and rubble, and the presence of small tide pools with standing water (Table 1). High levels of fine sedimentation on substrates are not conducive to coral recruitment and growth (Rogers 1990; Rogers *et al.* 1994). It is postulated that predominant north-westerly tidal movement drives local cross-shore transport of finer sediment from north to south.

The coral species and lifeforms in these extreme conditions are characteristic of those that occupy sheltered reefs with high sedimentation and turbidity (Rosen 1971; Chappel 1980; Done 1982). Species with massive, submassive and encrusting life forms are more tolerant to thermal stress (Marshall & Baird 2000), while convex and steep sided growth morphologies facilitate sediment flow off surfaces (Lasker 1980; Rogers 1990). The most diverse species were from the Merulinidae, which are represented by species that are tolerant of thermal stress (Depczynski *et al.* 2013; Schoepf *et al.* 2015) and well defined association with turbid water and lengthy periods of emersion. Corals with large polyps, like most species from the Merulinidae are efficient at removing sediment (Marshall & Orr 1931), and species such as *Dipastrea* spp., *Goniastrea* spp., *Lobophyllia hemphrichii*, *Astrea curta*, *Platygyra* spp., and *Symphyllia recta* can easily manipulate silt and fine sands (Stafford-Smith & Ormond 1992). Species of the genera *Coelastrea* and *Goniastrea*, for example, are often found in conditions where corals may not be expected to survive (Veron 1986) and are dominant in high latitude reefs in waters of high nutrients and turbidity (Thomson & Frisch 2010). In the Arabian Gulf region, the Merulinidae has a greater representation of species compared to other coral families such as Acroporidae which has greater species diversity in the Indo-Pacific region (Coles 1993; Foster & Foster 2013).

Species of the Acroporidae were scarce, with only two species represented at very low coral cover. The Acroporidae is reported to occur in other regions with high thermal stress (e.g. Craig *et al.* 2001; Bauman *et al.* 2013b; Dandan *et al.* 2015; Schoepf *et al.* 2015). At Nightcliff Reef it is probably further limited by the high turbidity as it is a poor sediment rejector with small polyps less than 2 mm diameter (Stafford-Smith & Ormond, 1992). However, a small-polyped species, *Porites* cf. *nigrescens* (Poritidae) was abundant on the middle and outer reef zones. This species has also been recorded as abundant amongst a coral community surveyed at Cobourg Peninsula, Northern Territory (Billyard 1995) and to the west on macro-tidal reef flats and lagoons off Sunday Island at the mouth of King Sound, Western Australia (Dobson 1999). In further regions, *Porities nigrescens* is reported as common on turbid fringing reefs to the south of Saudi Arabia (Jeddah to Jizan), but disappears to the north in clearer waters (Sheppard 1985). The species is regarded as a shallow water reef builder in environments with low light intensities and low wave energy (Cabioch *et al.* 1999). In northern Australia it appears to have adopted a similar niche to that occupied by *Montipora digitata* that is common on more sedimented



inshore reefs of eastern Australia, such as the reef flats of Magnetic Island, Queensland (Bull 1982; Mapstone *et al.* 1992).

Sediments for the upper and middle reef flat zones across the majority of localities were predominantly muds and silts on rock with intervening patches of muddy sand. Benthic algae were both more abundant and more diversely represented in these zones and collectively contributed to 38 species (Table 1). The brown foliose species *Padina australis* dominated the upper reef flat zone. Other conspicuous bladed foliose algae included Sargassaceae holdfasts (probably *Sargassopsis decurrens*), *Splatoglossum asperum* and *Dictyopteris* sp. Considerable turf forming algae were also present on the upper reef flat zone, and dominated in the middle reef flat zone with *Amphiroa fragilissima*, *Portieria hornemanni*, *Tricleocarpa fragilis*, *Acanthophora specifera*, *Gracilaria salicornia* and *Anadyomene plicata* visually conspicuous and varying in relative abundance across localities.

The lower reef flat zone, which was dominated by corals, exhibited minimal algal cover (0.03–0.05% mean cover, Table 1). There was no evidence of competition from algae with corals as described in eastern Australia (Morrissey 1980), the Arabian Gulf (Sheppard *et al.* 1992) and the Red Sea (Loya 1977b). It appears corals have a competitive advantage in this lower reef flat zone due to the greater water depth and ability to cope with the high turbidity which results in lower light availability to benthic algae. In the shallower upper tidal zones, longer and more frequent emersion times exclude most coral species, and this allows rapid colonisation of benthic algae which also gain improved light attenuation for photosynthesis.

The algal species recorded at Nightcliff Reef have been widely reported across the Indo-Pacific, from Tanzania (Jassund, 1976) to eastern Australia (Morrissey 1980; Ngan & Price 1980). The abundance of benthic algal species at Nightcliff Reef is likely to be seasonal with species exhibiting variable growth and dominance between wet and dry seasons (e.g. Benayahu & Loya 1977b; Lawson & John 1987; Vuki & Price 1994). The dominance of fleshy brown algae in the upper reef flat zone, with the co-occurrence of turfing algae is similar to the reef flat zonation described in the Gulf of Eilat, Red Sea (Benayahu & Loya 1977a, 1977b).

A notable observation was the limited representation of erect coralline algae from the genus *Halimeda* which is an important contributor to calcium carbonate accretion and grows abundantly on reef flats in eastern Australia (Morrissey 1980), Guam (Merten 1971) and the Gulf of Mannar, India (Rao 1972). However, in similar extreme environmental conditions such as the Arabian Gulf region, *Halimeda* is not recognised as a major component of inshore reefs (Sheppard 1985; Sheppard *et al.* 1992).

The coral species of Darwin Harbour offer valuable insights to physiological and evolutionary adaptations to persist in the most extreme environmental conditions. It is evident from this study and comparative investigations elsewhere that species with large polyps, thick skeletal tissue and massive or submassive lifeform strategies are amongst the most successful to surviving high sedimentation, nutrients and water temperatures.

---

The close proximity of Nightcliff Reef to coastal development, land-based sources of pollution and marine development combined with the changing climate pose future conservation challenges for this reef which is currently deemed at high risk to integrated local threatening processes (Burke *et al.* 2011).

## Acknowledgments

I am grateful to the Parks and Wildlife Commission of the Northern Territory for funding, and support from Michael Guinea, Ron Billyard, Ian Fulton, Belinda Oliver, Ian Price and John Collins. This research was conducted towards partial fulfillment of a Bachelor of Honours degree at the Charles Darwin University.

## References

- Australian Bureau of Meteorology (2013) *Record sea surface temperatures*. <<http://www.bom.gov.au/climate/enso/history/ln-2010-12/SST-records.shtml>> (accessed 15 November 2013).
- Bauman A.G., Baird A.H. and Cavalcante G.H. (2011) Coral reproduction in the world's warmest reefs: southern Persian Gulf (Dubai, United Arab Emirates). *Coral Reefs* 30, 405–413.
- Bauman A.G., Pratchett M.S., Baird A.H., Riegl B., Heron S.F. and Feary D.A. (2013a) Variation in the size structure of corals is related to environmental extremes in the Persian Gulf. *Marine Environmental Research* 84, 43–50.
- Bauman A.G., Feary D.A., Heron S.F., Pratchett M.S. and Burt J.A. (2013b) Multiple environmental factors influence the spatial distribution and structure of reef communities in the northeastern Arabian Peninsula. *Marine Pollution Bulletin* 72, 302–312.
- Benayahu Y. and Loya Y. (1977a) Space partitioning by stony corals, soft corals and benthic algae on the coral reefs of the northern Gulf of Eilat (Red Sea). *Helgolander Wissenschaftliche Meeresuntersuchungen* 30, 362–382.
- Benayahu Y. and Loya Y. (1977b). Seasonal occurrence of benthic-algae communities and grazing regulation by sea urchins at the coral reefs of Eilat, Red Sea. In: *Proceedings of Third International Coral Reef Symposium Volume 1: Biology* (ed. Taylor D.L.), pp. 383–390. Rosenstiel School of Marine and Atmospheric Science, Miami, Florida.
- Billyard R.W. (ed) (1995) *The Cultural and Ecological Significance of Popham Creek, Cobourg Peninsula*. Parks and Wildlife Commission of the Northern Territory, Darwin, Australia.
- Budd A.F., Fukami H., Smith N.D. and Knowlton N. (2012) Taxonomic classification of the reef coral family Mussidae (Cnidaria: Anthozoa: Scleractinia). *Zoological Journal of the Linnean Society* 166, 465–529.
- Burke L. M., Reynter K., Spalding M. and Perry, A. (2011) *Reefs at Risk Revisited*. World Resources Institute, Washington, DC.
- Cabioch G., Camoin G. and Montaggioni L. (1999) Postglacial growth history of a French Polynesian barrier reef tract, Tahiti, central Pacific. *Sedimentology* 46, 985–1000.
- Chappell J. (1980) Coral morphology, diversity and reef growth. *Nature* 286, 249–252.
- Chiappone M. and Sullivan K.M. (1991) A comparison of line transect versus linear percentage sampling for evaluating stony coral (Scleractinia and Milleporina) community similarity and area coverage on reefs of the central Bahamas. *Coral Reefs* 10, 139–154.
- Coles S. L. (1997) Reef corals occurring in a highly fluctuating temperature environment at Fahal Island, Gulf of Oman (Indian Ocean). *Coral Reefs* 16, 269–272.
- Coles S. L. (2003) Coral species diversity and environmental factors in the Arabian Gulf and the Gulf of Oman: a comparison to the Indo-Pacific region. *Atoll Research Bulletin* 507, 1–19.

- Cribb A.B. (1983) *Marine Algae of the Southern Great Barrier Reef. Part 1. Rhodophyta*. Australian Coral Reef Society Handbook No. 2, Brisbane.
- Cribb A.B. and Cribb J. W. (1985) *Plant Life of the Great Barrier Reef and Adjacent Shores*. University of Queensland Press, St. Lucia, Australia.
- Craig P., Birkeland C. and Belliveau S. (2001) High temperatures tolerated by a diverse assemblage of shallow-water corals in American Samoa. *Coral Reefs* 20, 185–189.
- Dandan, S.S., Falter, J.L., Lowe, R.J. and McCulloch, M.T. (2015) Resilience of coral calcification to extreme temperature variations in the Kimberley region, northwest Australia. *Coral Reefs* 34, 1151–1163.
- Depczynski M., Gilmour J. P., Ridgway T., Barnes H. *et al.* (2013) Bleaching, coral mortality and subsequent survivorship on a West Australian fringing reef *Coral Reefs* 32, 233–238.
- Dobson G. (1999) *Assessing and limiting predation and other sources of mortality in reseeded Trochus niloticus*. Unpublished MSc Thesis. Northern Territory University, Darwin.
- Done T.J. (1982) Patterns in the distribution of coral communities across the central Great Barrier Reef. *Coral Reefs* 1, 95–107.
- English S., Wilkinson C. and Baker V. (eds.) (1994) *Survey Manual for Tropical Marine Resources*. Australian Institute of Marine Science, Townsville.
- Foster K.A. and Foster G. (2013) Demography and population dynamics of massive coral communities in adjacent high latitude regions (United Arab Emirates). *PLoS ONE* 8, e71049.
- Gilmour J.P., Cooper T.F., Fabricius K.E. and Smith L.D. (2006) *Early Warning Indicators of Change in the Condition of Corals and Coral Communities in Response to Key Anthropogenic Stressors in the Pilbara, Western Australia*. Australian Institute of Marine Science, Western Australia.
- Glynn P.W. (1993) Coral reef bleaching: ecological perspectives. *Coral Reefs* 12, 1–17.
- Guiry M.D. and Guiry G.M. (2013) *AlgaeBase*. World-wide electronic publication, National University of Ireland, Galway. <<http://www.algaebase.org>> (accessed 12 December 2015).
- Hooper J.N.A. (1987). Structural features of the benthic community of East Point Fish Reserve. A comparative study between oceanic, nearshore and inshore reefs of Northwest Australia. In: *Proceedings of a workshop on research and management held in Darwin, 2–3 September, 1987* (eds Larson H.K., Michie M.G. and Hanley J.R.), pp. 215–225. Australian National University, North Australia Research Unit, Darwin.
- Huang D., Benzoni F., Fukami H., Knowlton N., Smith N. D. and Budd A. F. (2014) Taxonomic classification of the reef coral families Merulinidae, Montastraeidae, and Diploastraeidae (Cnidaria: Anthozoa: Scleractinia). *Zoological Journal of the Linnean Society* 171, 277–355.
- Hughes T. P., Baird A. H., Bellwood D. R., Card M. *et al.* (2003). Climate change, human impacts, and the resilience of coral reefs. *Science* 301, 929–933.
- Jaasund E. (1976) *Intertidal Seaweeds in Tanzania: A Field Guide*. University of Tromsø Press, Norway.
- Lasker H.R. (1980). Sediment rejection by reef corals: the role of behaviour and morphology in *Montastrea cavernosa* (Linnaeus). *Journal of Experimental Marine Biology and Ecology* 47, 77–87.
- Lawson G.W. and John D.M. (1987) *The Marine Algae and Coastal Environment of Tropical West Africa*. J. Cramer Publisher, Berlin.
- Mapstone B.D., Choat J.H., Cumming R.L. and Oxley W.G. (1992) *The Fringing Reefs of Magnetic Island: Benthic Biota and Sedimentation – a Baseline Study*. Great Barrier Reef Marine Park Authority Research Publication No.13, Townsville.
- Marshall P.A. and Baird A.H. (2000) Bleaching of corals on the Great Barrier Reef: differential susceptibilities among taxa. *Coral Reefs* 19, 155–163
- Marshall S.M. and Orr A.P. (1931) Sedimentation on Low Isles Reefs and its relation to coral growth. *Scientific Report of the Great Barrier Reef Expedition* 3, 273–312.
-

- Merten M.J. (1971) Ecological observations on *Halimeda macroloba* Decaisne (Chlorophyta) on Guam. *Micronesia* 7, 27–44.
- Michie M.G. (1987) Distribution of Foraminifera in a macrotidal tropical estuary, Port Darwin, Northern Territory of Australia. *Australian Journal of Marine and Freshwater Research* 38, 249–259.
- Morrissey J. (1980) Community structure and zonation of macroalgae and hermatypic corals on a fringing reef flat of Magnetic Island Queensland, Australia. *Aquatic Botany* 8, 91–139.
- Ngan Y. and Price L.R. (1980) Distribution of intertidal benthic algae in the vicinity of Townsville, Tropical Australia. *Australian Journal of Marine and Freshwater Research* 31, 175–191.
- Price I.R. and Scott F.J. (1992) *The Turf Algal Flora of the Great Barrier Reef. Part 1 Rhodophyta*. James Cook University of North Queensland Press, Townsville.
- Rao M. (1972) Coral reef flora of the Gulf of Mannar and Palk Bay. In *Proceedings of the 1st International Symposium on Corals and Coral Reefs* (eds Mukundan M. and Gopinadha Pillai C.S.), pp. 217–230. Marine Biological Association of India, India.
- Riegl B. (1999) Corals in a non-reef setting in the southern Arabian Gulf (Dubai, UAE): fauna and community structure in response to recurring mass mortality. *Coral Reefs* 18, 63–73.
- Riegl B. and Purkis S. (eds) (2012) *Coral Reefs of the Gulf: Adaptation to Climatic Extremes*. Coral Reefs of the World Volume 3, Springer, Netherlands.
- Rogers C.S. (1990) Responses of coral reefs and reef organisms to sedimentation. *Marine Ecology Progress Series* 62, 185–202.
- Rogers C.S., Fitz H.C., Gilnack M., Beets J. and Hardin J. (1984) Scleractinian coral recruitment patterns at Salt River Submarine Canyon. St. Croix, U.S.V.I. *Coral Reefs* 3, 69–76.
- Rosen B.R. (1971) Principal features of reef coral ecology in shallow water environments of Mahe, Seychelles. *Symposium of the Zoological Society of London* 28, 163–183.
- Schoepf V., Stat M., Falter J.L. and McCulloch M. (2015) Limits to thermal tolerance of corals adapted to a highly fluctuating, naturally extreme environment, *Science Report* 5, Article 17639.
- Semeniuk V. (1985) Mangrove environments of Port Darwin, Northern Territory: the physical framework and habitats. *Journal of the Royal Society of Western Australia* 67, 81–97.
- Sheppard C. and Sheppard A. (1991) Corals and coral communities of Arabia. *Fauna of Arabia* 12, 3–17.
- Sheppard C.R.C., Price A. and Roberts C. (1992) *Marine Ecology of the Arabian Region. Patterns and Processes in Extreme Tropical Environments*. Academic Press, London.
- Stafford-Smith M.G. and Ormond R.F.G. (1992) Sediment-rejection mechanisms of 42 species of Australian scleractinian corals. *Australian Journal of Marine and Freshwater Research* 43, 683–705.
- Sullivan K.M. and Chiappone, M. (1992) A comparison of belt quadrat and species presence/absence sampling of stony coral (Scleractinia and Milleporina) and sponges for evaluating species patterning on patch reefs of the central Bahamas. *Bulletin of Marine Science* 50, 464–488.
- Veron J.E.N. (1986) *Corals of Australia and the Indo-Pacific*. Angus and Robertson, Australia.
- Veron J.E.N. and Pichon M. (1976) *Scleractinia of Eastern Australia. Part I. Families Thamnasteriidae, Astrocoeniidae, Pocilloporidae*. AIMS Monograph Series 1, Australian Institute of Marine Science, Townsville.
- Veron J.E.N. and Pichon M. (1980) *Scleractinia of Eastern Australia. Part III. Families Agaraciidae, Siderastreidae, Fungiidae, Oculinidae, Merulinidae, Mussidae, Pectiniidae, Caryophylliidae, Dendrophylliidae*. AIMS Monograph Series 4, Australian Institute of Marine Science, Townsville.
- Veron J.E.N. and Pichon M. (1982) *Scleractinia of Eastern Australia. Part IV. Families Poritidae*. AIMS Monograph Series 5, Australian Institute of Marine Science, Townsville.
-

- Veron J.E.N., Pichon M. and Wijsman-Best M. (1977) *Scleractinia of Eastern Australia. Part II. Families Faviidae, Trachyphyllidae*. AIMS Monograph Series 3, Australian Institute of Marine Science, Townsville.
- Veron J.E.N. and Wallace C.C. (1984) *Scleractinia of Eastern Australia. Part V. Family Acroporidae*. AIMS Monograph Series 6, Australian Institute of Marine Science, Townsville.
- Vuki V.C. and Price I.R. (1994) Seasonal changes in the *Sargassum* populations on a fringing reef, Magnetic Island, Great Barrier Reef Region, Australia. *Aquatic Botany* 48, 153–166.
- Wolstenholme J., Dinesen Z.D. and Alderslade P. (1997) Hard corals of the Darwin region, Northern Territory, Australia. In: *Proceedings of the Sixth International Marine Biological Workshop: The Marine Flora and Fauna of Darwin Harbour, Northern Territory, Australia* (ed. Hanley R), pp. 381–398. Museum and Art Gallery of the Northern Territory and the Australian Marine Sciences Association, Darwin.
- WoRMS Editorial Board (2015) *World Register of Marine Species*. <<http://www.marinespecies.org>> (accessed 24 November 2015).
- Wynne M.J. (2011) The benthic marine algae of the tropical and subtropical Western Atlantic: changes in our understanding in the last half century. *Algae* 26, 109–140.
-